

## Ceftiofur as an Intrauterine Therapy in Treating Endometritis in Cows

Sarah Jasim Abdulameer<sup>1</sup>, N.Y.Khudhair<sup>2</sup>

<sup>1</sup>Microbiology Department, College of Vet. Medicine, University of Diyala, Iraq

<sup>2</sup>Surgery & Obstetrics Department, College of Vet. Medicine, University of Diyala, Iraq

\*Corresponding Author: E.mail: [sarah.j@uodiyala.edu.iq](mailto:sarah.j@uodiyala.edu.iq)

### Abstract

#### Aims:

Recognition the efficacy of ceftiofur antibiotic as intrauterine therapy in treating clinical endometritis and the range of its influence on the fertility of treated cows.

#### Methods:

Twenty multiparous cows suffering from clinical endometritis(CE) came to the veterinary clinic exist in the college and veterinary clinics located in Diyala province. Rectal palpation was made to diagnose the clinical endometritis, all these cows were treated by intra-uterine injection of ceftiofur during estrus phase. After the coming of all treated cows to estrus they were inseminated artificially, and then the rectal palpation was carried out to confirm the pregnancy in these treated cows.

#### Results:

The number of multiparous cows had clinical endometritis and treated with ceftiofur intra-uterine which were 20. All these cows came estrus within 18-23 days post treatment, and inseminated artificially. Eight cows were return to estrus within 20-22 days after last insemination and re-inseminated again. After 55-60 days of last insemination all cows were examined by rectal palpation and being pregnant.

**Key words:** Cows, Ceftiofur, Clinical endometritis , Fertility

### Introduction

Clinical endometritis (CE) is the inflammation of the endometrial layer of the uterus without an association to systemic signs (1), (2). An increased rectal temperature may or may not be present depending upon the extent of the inflammatory process and stress level of the cow at the time of the examination. CE is characterized by muco-purulent and purulent vaginal discharge around 20 to 40 days postpartum (3). CE has been known as the presence of muco-purulent uterine discharge (>50 % purulent material)

after 26 days in milk or a diameter of cervix greater than 7.5 cm (by rectal examination (3, 4). Involution of the uterus begins immediately post-partum and the diameter of the uterine horns should be less than 3-4 cm and the cervical diameter should measure <5 cm by 40 days in milk (5). The diagnosis of CE relies on appropriate visual characterization of the uterine discharges noted at 21 days postpartum. Vaginoscopy is a practical method used for the diagnosis of clinical endometritis(CE) (uterine discharge)(6,7)scored using a 0-3 scale

(0 = normal uterine discharge, 1 = flakes of purulent exudate in the uterine discharge, 2 = <50% of the uterine discharge is made up of purulent exudate, 3 = uterine discharge mixed with >50% purulent exudate(8)). Palpation of the uterus per rectum to estimate the uterine diameter and the presence of intraluminal fluid is commonly the sole method used for diagnosing CE. Palpation per rectum combined with vaginoscopy results in a more accurate method of diagnosis than palpation alone (9,11).

The incidence rate for endometritis in cows varies considerably among studies, perhaps owing to different methods used to classify the disease. A review of the literature shows that the percentage of cows affected with endometritis can be as high as 50% at 40–60 days in milk (10). Additional studies have reported a range of 16.9% to 34% prevalence of CE for cows examined between 21 and 28 days postpartum(12,8,3). however, based on the method used (cytology, biopsy, vaginoscopy, or rectal palpation) for diagnosis, the incidence of endometritis likely lies between 7.5% to 34% (10). A diagnosis of endometritis based on unspecified criteria has been reported to be 7.8–13.8 % (13, 14). An 18% incidence of endometritis was reported when the diagnosis was made using palpation per rectum. A mini-review published in 2009 summarized the rates and diagnostic criteria for endometritis (4).

Impact of clinical endometritis on the cow's fertility

Under normal circumstances, the genital tract of dairy cows should have recovered and able to establish and maintain a new pregnancy 45 days post-calving (4). Cows that exhibit CE take approximately 27% longer to get pregnant than normal cows (7) and are 1.7 times more likely to be culled due

to reproductive failure (10). Cows with CE had a 17% reduction in pregnancy rate relative to cows with normal uterine discharge (7). Cows with endometritis have a calving interval that is 30 days longer than cows without endometritis (15). Endometritis is responsible for increased culling of cows due to reproductive failure (4). Purulent or mucopurulent discharge was associated with significantly lower pregnancy and first service conception rates (6).

Ceftiofur is a third-generation cephalosporin, an important class of antimicrobial drugs used in veterinary medicine (16). Furthermore, intrauterine infusion of ceftiofur hydrochloride positively affected uterine health in dairy cows (17). Ceftiofur has high effectiveness against all gram-positive and gram-negative pathogenic microorganisms (18). In the liver, it is converted (by hydrolysis process) into desfuroyl-ceftiofur-acetamide (**DCA**), which is an active metabolite that is protected from fast renal remove through protein binding (17).

There are few previous studies about using ceftiofur for treatment clinical endometritis in dairy cows, so this study will test this drug for treatment the cows that suffered from endometritis.

### Materials and Methods

This study was conducted in college of Veterinary Medicine in Diyala governorate in a period extends from 15/2/2024 until 1/9/2024, and included twenty multiparous cows suffering from clinical endometritis(CE), and came to the veterinary clinic exist in the college and other veterinary clinics located in the province. After the taking the case history of these cows, rectal palpation was made to diagnose the clinical endometritis, which

characterized by mucopurulent and purulent vaginal discharge and with a cervical diameter greater than 7.5 cm. after the diagnosis of (CE), all these cows were treated by intra-uterine injection of one thousands mgs (20 ml) of ceftiofur (CEFTIOFUR-VS, ((VETSINTEZ)) LLC, Kharkiv-Ukraine)(each 1ml contains 50 mg Ceftiofur hydrochloride), during estrus phase. All treated cows came to estrus within 18-23 days post treatment and were inseminated artificially, and after 20-22 days of insemination 8 cows were return and re-inseminated. After 55-60 days of late, insemination the pregnancy diagnosis was done rectally to confirm the gestation.

Statistical Analysis

The data were analyzed statistically by using Chi-square to significant compare the results at (0.05) in this study (19).

### Results

As shown in table (1) the number of multiparous cows had clinical endometritis and treated with ceftiofur intra-uterine which were 20. All these cows came to estrus within 18-23 days after treatment, and had been inseminated, 8 cows were return to estrus within 20-22 days after last insemination and re-inseminated again and after 55-60 days of last insemination all cows were examined by rectal palpation and being pregnant.

**Table 1:** Distribution of sample study according to onset of estrus after treatment with ceftiofur.

	No and Period	No and Period	P-value
Cows	20 cows treated with ceftiofur	12 cows not return to estrus (60.00%)	0.0349 *
		8 cows return to estrus (40.00%)	
Time of onset of estrus after treatment	18- 23 days	20 - 22 days	---
Rectal palpation after last insemination	55- 60 days	20 (100%) pregnant	---
* ( $P \leq 0.05$ ).			

### Discussion

The findings of the current study showed that all twenty cows suffering from endometritis responded to the treatment by ceftiofur intrauterine therapy and came to estrus after 18-23 days after treatment and inseminated by using artificial insemination (AI). twelve (60.00%) cows not return to estrus secondly, whilst eight (40.00%) cows return to estrus within 20-22 days after last insemination and re-

inseminated again and all inseminated treated cows being pregnant after 55-60 days of artificial insemination. This reveals that the cefitiofur was efficient to treat clinical endometritis in treated cows in this study.

These results are in accord with many studies indicated that cefitiofur has good efficacy in treating uterine disease such the metritis and clinical endometritis and has good role in

enhancement of uterine health in the cows infected with uterine disease(20,21,22). However another studies disagreed with these results and demonstrated that ceftiofur hydrochloride has no effect on the reproductive performance of infected cows with endometritis(17,23). (Coto and Lucy, 2018) believed that ceftiofur was effective only in cows that did not suffer retained placenta and with two or more parturitions. Ceftiofur was efficient in removing the most causative agents of clinical endometritis more than its ability to enhance reproductive performance of treated cows suffering endometritis, and the ability of this antibiotic may be potentiated if is associated with the usage of uterine tonic hormones like oxytocin or prostaglandinF2 $\alpha$ (24). Many previous studies that had employed ceftiofur in treating of uterine diseases like clinical endometritis and puerperal metritis recommended using this antibiotic parentally either intramuscularly or subcutaneously; as this antibiotic was highly effective in treating uterine disease in cows if it was used by this method of administration (21, 22, 26, 27). The dose of ceftiofur used in this study was relatively high (1000 mgs) and led to good percentage of recovery which was 60%. It was well known that the highly dose of antibiotic or drug would reinforce the response to the antibiotic by the treated animals and this resulted in enhancement of reproductive performance of the involved cows in this study which reflected on the overall pregnancy rate which, was high( 27, 28, 29).

#### Conclusion:

From these results we can concluded that antibiotic ceftiofur was highly efficient in treating clinical

endometritis in cows intra-uterine, with better results if it is used with highly doses.

#### References

- 1- Sheldon, I.M.; Lewis, G.S.; LeBlanc, S., et al. (2006) Defining postpartum uterine disease in cattle. *Theriogenology*; 65:1516-1530.
- 2- Barlund, C.S; Carruthers, T.D.; Waldner, C.L., et al. (2008) A comparison of diagnostic techniques for postpartum endometritis in dairy cattle. *Theriogenology*; 69:714-723.
- 3- Mahmood, M.Y.; Rasheed, Y.M.; Ghareeb, S.H. & Talib, N.T. (2024) Subclinical endometritis: Fungal identification and treatment of conception failure in cows. *DJVS*; 2(3): 67-74.
- 4- Sheldon, I.M.; Cronin, J.; Goetze, L. et al. (2009) Defining Postpartum Uterine Disease and the Mechanisms of Infection and Immunity in the Female Reproductive Tract in Cattle. *Biol Reprod*; 123:837-845.
- 5- Morrow, D.A.; Roberts, S.J.; McEntee, K.; Gray, H.G.(1966) Postpartum ovarian activity and uterine involution in dairy cattle. *American Veterinary Medical Association*; 149:1596-1609.
- 6- Runciman, D.J.; Anderson, G.A.; Malmo, J. et al. (2008) Use of postpartum vaginoscopic (visual vaginal) examination of dairy cows for the diagnosis of endometritis and the association of endometritis with reduced reproductive performance. *Aust Vet J*; 86:205-213.
- 7- LeBlanc, S.J.; Duffield, T.F.; Leslie, K.E. et al. ( 2002) Defining and diagnosing postpartum clinical endometritis and its impact on reproductive performance in dairy cows. *J Dairy Sci*; 85:2223-2236.
- 8- Williams, E.J.; Fischer, D.P.; Pfeiffer, D.U. et al. (2005) Clinical

- evaluation of postpartum vaginal mucus reflects uterine bacterial infection and the immune response in cattle. *Theriogenology*; 63:102-117.
- 9- Miller, H.V.; Kimsey, P.B.; Kendrick, J.W., et. al. (1980) Endometritis of dairy cattle: diagnosis, treatment, and fertility. *Bov Prac.*; 15: 13-23.
  - 10- Gilbert, R.O.; Schwark, W.S.(1992) Pharmacologic considerations in the management of peripartum conditions in the cow. *Vet Clin North Am Food Anim Pract*; 8:29-56.
  - 11- LeBlanc, S.J.; Duffield T.F.; Leslie, K.E., et al. (2002). The effect of treatment of clinical endometritis on reproductive performance in dairy cows. *J. Dairy Sci*;85: 2237-2249.
  - 12- Heuwieser, W., Tenhagen, B.A., Tischer, M. et al. (2000) Effect of three programs for the treatment of endometritis on the reproductive performance of a dairy herd. *Vet Rec*; 146:338-341.
  - 13- Bartlett, P. (1986) Metritis complex in Michigan Holstein-Friesian cattle: incidence, descriptive epidemiology and estimated economic impact. *Prev Vet Med*; 4:235-248.
  - 14- Mohammed, S.N. (2024) Bacteriological finding, Vaginal discharges, and endometrial cytology for Endometritis Detection in Postpartum Buffaloes. *DJVS*; 2(3): 31-49.
  - 15- Borsberry, S.; Dobson, H. (1989) Periparturient diseases and their effect on reproductive performance in five dairy herds. *Vet Rec*; 124:217-219.
  - 16- Hornish, R. E. and Kotarski, S. F. (2002) Cephalosporins in veterinary medicine – Ceftiofur use in food animals. *Curr. Top. Med. Chem.*, 2:717–731.
  - 17- Witte, T. S.; Iwersen, M.; Kaufmann, T.; Scherpenisse, P.; Bergwerff, A.A. and Heuwieser, W. (2011) Determination of ceftiofur derivatives in serum, endometrial tissue, and lochia in puerperal dairy cows after subcutaneous administration of ceftiofur crystalline free acid. *J.Dairy.Sci.*, 94:284–290.
  - 18- Haggett, E. F. and W. D. Wilson. (2008) Overview of the use of antimicrobials for treatment of bacterial infections in horses. *Equine Vet. Educ.* 20:433–448.
  - 19- SAS, (2018) Statistical Analysis System, User's Guide. Statistical. Version 9.6<sup>th</sup> ed. SAS. Inst. Inc. Cary. N.C. USA.
  - 20- Coto, R.M. and Lucy, M.C.(2018) Uterine inflammation affects the reproductive performance of dairy cows: A review. *Agron. Mesoam.* 29(2):449-468.
  - 21- El-Khadrawy, H.H., Ahmed, W.M., Zaabal, M.M., Hanafi, E.M.(2015) Strategies for Diagnosis and Treatment of Uterine Infection in Bovines. *Global Veterinaria* 15 (1): 98-105.
  - 22- Galvao, K.N., Greco, L.F., Vilela, J.M., Sa Filho, M.F., and Santos, J.E.P.(2009) effect of intrauterine infusion of ceftiofur on uterine health and fertility in dairy cows. *J. Dairy Sci.* 92 :1532–1542.
  - 23- Fard, A.A.G., Batavani, R.A. and Amanalou, H. (2017) Investigation the Efficiency Ceftiofur Hydrochloride in Prevention of Uterine Postpartum Infections and Improvement the Fertility Parameters in Holstein Dairy Cows Affected by Retained Fetal Membrane and Dystocia. *International Journal of Advanced Biological and Biomedical Research* 5(2) (65–68).
  - 24- Kacar, C. and Kaya, S. (2014) Uterine Infections in Cows and Effect on Reproductive Performance. *Kafkas Univ Vet Fak Derg.*, 20(6): 975-982.



- 25- Galvao, K.N. (2012) Postpartum uterine diseases in dairy cows. Anim Reprod, v.9, n.3, p.290-296 .
- 26- Brick, T.A. (2011) Impact of Intrauterine Dextrose Therapy on Reproductive Performance of Lactating Dairy Cows Diagnosed with Clinical Endometritis Following a Randomized Clinical Trial. A thesis, Ohio University, USA.
- 27- Pyorala, S., J. Taponen, and T. Katila. (2014) Use of antimicrobials in the treatment of reproductive diseases in cattle and horses. Reprod. Domest. Anim. 49:16-26.
- 28- Noakes, D.E., Parkinson, T.J. and England, G.C.W. (2009) Veterinary Reproduction and Obstetrics. Ninth edition, Saunders Elsevier, U.K.
- 28- Hafez, B. and Hafez, E.S.E.(2000) Reproduction in Farm Animals. Seven edition, Lipincott Williams and Wilkins, USA.
- 29- Hafez, B. and Hafez, E.S.E.(2000) Reproduction in Farm Animals. Seven edition, Lipincott Williams and Wilkins, USA.